

WARNING
KEEP OUT OF REACH OF CHILDREN
READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

KENSTRIKE

HERBICIDE

ACTIVE CONSTITUENT: 800 g/kg FLUMETSULAM in the form of a water dispersible granule

GROUP **2** HERBICIDE

For post emergence control of broadleaf weeds in clover, lucerne, chicory, new and mature pasture and maize



NET CONTENTS: 500 g

Registered pursuant to the ACVM Act 1997, No. P009082
See www.foodsafety.govt.nz for registration conditions
Approved pursuant to the HSNO Act 1996, Approval No. HSR101014
See www.epa.govt.nz for approval controls

Registrant:
Kenso Corporation (M) Sdn Bhd
Lot 29, Jalan Sungai Pinang 5/4, Taman Perindustrian
Pulau Indah, 42920 Pelabuhan Klang, Selangor, Malaysia.

NZ Agent:
Kenso NZ Limited
2 Bond Crescent, Forrest Hill, Auckland 0620 New Zealand.
Ph: 0800 536 766

KENSTRIKE HERBICIDE

PERSONNEL QUALIFICATION & EXPERIENCE

This product must be under the control of an appropriately trained and experienced applicator when it is applied in a wide dispersive manner, or used by a commercial contractor. This product must not be used for any purpose or in any manner, contrary to this label unless authorised under appropriate legislation.

HAZARD CLASSIFICATION

Hazardous to the aquatic environment acute Category 1, Hazardous to the aquatic environment chronic Category 1, Hazardous to soil organisms.

ECOTOXIC

Very toxic to aquatic organisms with long lasting effects. Avoid contamination of any water supply with product or empty container.

Selective herbicide — very toxic to some plant species.

Beware: Spray drift hazard. Apply this product carefully. Spray drift may cause serious damage to other desirable plants. Do not allow drift to occur outside the target area.

HANDLING PRECAUTIONS

STORAGE

Keep out of reach of children. Store in original container, tightly closed and in a locked, dry, cool area away from foodstuffs, fertilisers and seeds. Store in accordance with NZS 8409 Management of Agrichemicals. Stores containing 100 kg of Kenstrike Herbicide require signage and more than 100 kg require emergency response plans.

DISPOSAL

Triple rinse empty container and add rinsate to spray tank. Recycle empty container through Agrecovery (0800 247 326, www.agrecovery.co.nz). Otherwise, crush and bury in a suitable landfill. Dispose of product only by using according to the label, or at an appropriate landfill.

SPILLAGE

Wear appropriate protective clothing and prevent material from entering waterways. Contain liquid spills, and absorb with inert material. Sweep up solids without creating dust. Dispose of waste safely in an approved landfill.

Batch No.:

Date of Manufacture:

FIRST AID

If swallowed, DO NOT induce vomiting. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. If splashed in eyes, wash out immediately with water. If inhaled, move the victim to fresh air immediately. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately.

SAFETY DATA SHEET

A Safety Data Sheet is available from the Kenso NZ website: www.kenso.co.nz

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.



EMERGENCY RESPONSE
Ring 0800 CHEMCALL, or
0800 243 622 (24 hours)

Product Of China



PERSONAL PROTECTION

Avoid skin and eye contact and inhalation of spray mist. When mixing or applying wear appropriate eye protection, waterproof gloves, boots and overalls. Wash hands and exposed skin before meals and after work.

GENERAL INFORMATION

Kenstrike Herbicide will control or suppress broadleaf weeds in crops or pasture while selectivity to clovers and chicory is preserved.

APPLICATION

Make sure that spray equipment is correctly calibrated. Ensure complete spray coverage for best results. Do not overspray.

Aerial and ground broadcast application: Apply Kenstrike Herbicide in 100 - 300 litres of water per hectare. Use the higher water rate where weeds are dense.

CDA application: Use the large (250 - 300 micron) droplet size and apply not less than 30 litres of water per hectare. Trials indicate that CDA application may result in less effective weed control than conventional boom spraying.

DIRECTIONS FOR USE

Apply to weeds in active growth. Do not apply in cold, wet conditions. Always apply Kenstrike Herbicide in mixture with a suitable spraying oil such as Koil at 500 ml/100 litres of water, unless it is specifically recommended not to do so.

WEEDS CONTROLLED	RATE g/ha	CRITICAL COMMENTS
Chickweed, Spurrey, Wild Radish, Hedge Mustard.	30	Apply before 4 th leaf.
Amaranthus (Redroot), Cleavers, Black Nightshade, Fathen, Shepherd's Purse, Mallow.	50	Apply before 4 th leaf.
Annual Buttercup, Creeping Yellow Cress, Yellow Gromwell.		Apply before flowering.
Larger weeds (above). Stinking Mayweed, Wireweed, Sorrel. Field Pansy, Henbit, Willow Weed, Oxeye Daisy. Giant Buttercup.	65	Apply to young seedlings. Apply to young seedlings. Suppression only. Apply before flowering.

Mature Pasture: Kenstrike Herbicide may be applied in either spring or autumn for buttercup control. Repeat spring treatments for two successive seasons will maximise control of giant buttercup.

Application Timing

CROP	CROP GROWTH STAGE
New pasture	Use after clover has developed two trifoliolate leaves.
New and established White Clover Seed Crops	Apply after the emergence of the 4 th trifoliolate leaf and before stolon elongation.
New and established Red Clover Seed Crops and Subterranean clover	Apply after the emergence of the 6 th trifoliolate leaf and before stolon elongation.
Established Lucerne	Apply before flowering.
Maize	Apply at the 4 th leaf (90mm plant height) stage.
New and established Chicory	Apply after the emergence of the 4 th leaf.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residues Levels for Agricultural Compounds.

WITHHOLDING PERIODS

Do not graze crops or treated pasture for 14 days after treatment.

COMPATIBILITY

The addition of Helion Herbicide or Pasture Star Herbicide to Kenstrike Herbicide will increase the range of broadleaf weeds controlled. Also compatible with most commonly used herbicides. Do not mix Kenstrike Herbicide with spraying oil when using with triazine herbicides on lucerne.

The use of spraying oil with Kenstrike Herbicide and Javelin Herbicide on pasture will increase clover damage and is not recommended unless the pasture has been well grazed before spraying.

EQUIPMENT

Apply using well maintained and accurately calibrated spray equipment. Clean equipment thoroughly after use.

CROP ROTATION

Do not plant susceptible crops until three months after application of Kenstrike Herbicide. Susceptible crops include all brassica species, sunflowers, lupins and beans. Do not allow spray to drift onto susceptible crops or land to be planted with susceptible crops.

ANIMAL SAFETY

Poisonous plants (e.g. including but not limited to giant buttercup and black nightshade) may become more palatable after spraying. If concerned that stock health may be affected, stock should be kept away from the treated area until plants have died down.

CLEANING SPRAY EQUIPMENT AFTER USE

Drain completely and flush system twice with clean water immediately after spraying. Discard rinse water on land already sprayed or on waste land away from desirable plants and water sources. If a susceptible crop (see above) is to be sprayed next, additional cleaning with detergent is recommended. In this case, after the initial flush with clean water, refill the spray tank adding an alkali detergent to make a 0.5 percent solution (e.g. — a low foam laundry detergent added at 500 g/100 litres of water). Flush through the boom and hoses, and allow to stand with agitation engaged for 15 minutes. Remove filters and nozzles and clean them separately. Drain the system and flush thoroughly with clean water.

RESISTANCE MANAGEMENT | | | | |-------|---|-----------| | GROUP | 2 | HERBICIDE | |-------|---|-----------|

Kenstrike Herbicide contains a GROUP 2 HERBICIDE belonging to the triazolopyrimidine sulfonanilide (sulfonamide) group of chemistry, with the inhibition of acetolactate synthase (ALS) mode of action. Naturally occurring biotypes resistant to sulfonanilide herbicides are known to exist. Resistance to these herbicides could develop from excessive use. To minimise the risk use strictly in accordance with label instructions. Refer to the NZCPR web site for more detailed information (www.nzpps.org/resistance/index.php).

TRACKING/RECORD KEEPING

Records of use, as described in NZS 8409 Management of Agrichemicals, must be kept if more than 3 kg of Kenstrike Herbicide is applied within 24 hours.

SHELF LIFE

When stored appropriately this product should show no significant degradation for 2 years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

UN NO: 3077



**ENVIRONMENTALLY HAZARDOUS
SUBSTANCE, SOLID, N.O.S.
(CONTAINS FLUMETSULAM)
PACKING GROUP III,
HAZCHEM 2X
HAZARD CLASS 9**

**In a transport emergency dial 111, Police or Fire Service.
For specialist advice in emergency only, call 0800 243 622 (24 hours)**